



# In Vitro Interactions of Echinocandins with Triazoles against Multidrug-Resistant *Candida auris*

Hamed Fakhim,<sup>a,b</sup>  Anuradha Chowdhary,<sup>c</sup> Anupam Prakash,<sup>c</sup> Afsane Vaezi,<sup>d</sup>  
 Eric Dannaoui,<sup>e</sup>  Jacques F. Meis,<sup>f,g</sup>  Hamid Badali<sup>h,i</sup>

Department of Medical Parasitology and Mycology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran<sup>a</sup>; Cellular and Molecular Research Center, Urmia University of Medical Sciences, Urmia, Iran<sup>b</sup>; Department of Medical Mycology, Vallabhbai Patel Chest Institute, University of Delhi, Delhi, India<sup>c</sup>; Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran<sup>d</sup>; Université Paris-Descartes, Faculté de Médecine, APHP, Hôpital Européen Georges Pompidou, Unité de Parasitologie-Mycologie, Service de Microbiologie, Paris, France<sup>e</sup>; Department of Medical Microbiology and Infectious Diseases, Canisius-Wilhelmina Hospital, Nijmegen, The Netherlands<sup>f</sup>; Centre of Expertise in Mycology Radboudumc/CWZ, Nijmegen, The Netherlands<sup>g</sup>; Department of Medical Mycology and Parasitology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran<sup>h</sup>; Pharmaceutical Sciences Research Center, Mazandaran University of Medical Sciences, Sari, Iran<sup>i</sup>

**ABSTRACT** We determined the *in vitro* interactions between echinocandins and azoles against 10 multidrug-resistant *Candida auris* strains by use of a microdilution checkerboard technique. Our results suggest synergistic interactions between micafungin and voriconazole with fractional inhibitory concentration index (FICI) values of 0.15 to 0.5, and we observed indifferent interactions when micafungin was combined with fluconazole (FICI, 0.62 to 1.5). Combinations of caspofungin with fluconazole or voriconazole exhibited indifferent interactions. No antagonism was observed for any combination.

**KEYWORDS** *in vitro* interactions, azoles, echinocandins, *Candida auris*

Candidiasis infection caused by uncommon *Candida* species has increased in recent years, particularly among immunocompromised patients (1). In the *Metschnikowia* clade, *Candida auris* causes various infections, ranging from superficial mucocutaneous candidiasis to severe bloodstream infections (2, 3). Remarkably, in recent years, multidrug-resistant *C. auris* has emerged in Asia, Africa, Europe, and the Americas, resulting in several cases of fungemia (3–14). Although European Society of Clinical Microbiology and Infectious Diseases (ESCMID) guidelines for the diagnosis and management of candidiasis recommend the use of azoles, polyenes, and echinocandins (15, 16), toxic effects of amphotericin B restrict its clinical application. In addition, resistance to azoles and echinocandins in *Candida* species has become a severe clinical challenge (17). Fungemia due to *C. auris* is associated with a high mortality rate and treatment failure, in addition to being potentially resistant to azoles, polyenes, and echinocandins (18–21). Thus, accurate identification of *C. auris* and *in vitro* antifungal susceptibility testing are highly recommended (22). Because of the limited available treatment choices and high rate of therapeutic failures, novel strategies are needed to improve patient outcomes (23). Combinations of echinocandins and azoles seem to be attractive treatment regimens, as both drug groups have different antifungal targets and modes of action. We therefore investigated the efficacy of echinocandins plus azoles against multidrug-resistant *C. auris* clinical isolates.

We studied 10 *C. auris* strains from patients with candidemia in tertiary care hospitals in Delhi, including fluconazole-resistant ( $n = 10$ ) and micafungin-resistant ( $n = 3$ ) isolates (according to non-species-specific *Candida* species breakpoints of >4

**Received** 19 May 2017 **Returned for modification** 15 July 2017 **Accepted** 20 August 2017

**Accepted manuscript posted online** 28 August 2017

**Citation** Fakhim H, Chowdhary A, Prakash A, Vaezi A, Dannaoui E, Meis JF, Badali H. 2017. *In vitro* interactions of echinocandins with triazoles against multidrug-resistant *Candida auris*. Antimicrob Agents Chemother 61:e01056-17. <https://doi.org/10.1128/AAC.01056-17>.

**Copyright** © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Hamid Badali, badali@yahoo.com.

**TABLE 1** *In vitro* interactions of caspofungin with fluconazole and voriconazole against *Candida auris*

Strain no.	CAS + FLU <sup>b</sup>				CAS + VRC <sup>b</sup>			
	MIC (μg/ml)				MIC (μg/ml)			
	CAS	FLU	CAS/FLU	FICI/INT	CAS	VRC	CAS/VRC	FICI/INT
VPCI 482/P/13 <sup>a</sup>	2	≥64	1/32	0.75/IND	2	2	1/0.5	0.75/IND
VPCI 1132/P/13 <sup>a</sup>	2	32	1/8	0.75/IND	2	0.5	1/0.063	0.62/IND
VPCI 1133/P/13 <sup>a</sup>	4	≥64	2/64	1/IND	4	1	2/0.25	0.75/IND
VPCI 265/P/14 <sup>a</sup>	4	32	2/32	1.5/IND	4	8	2/0.25	0.75/IND
VPCI 1510/P/14 <sup>a</sup>	0.5	32	0.5/32	2/IND	0.5	4	0.5/4	2/IND
VPCI 1514/P/14 <sup>a</sup>	1	≥64	0.5/32	0.75/IND	1	0.5	1/0.25	1.5/IND
VPCI 266/P/14 <sup>a</sup>	2	≥64	1/32	0.75/IND	2	0.5	1/0.25	1/IND
VPCI 267/P/14 <sup>a</sup>	2	32	1/8	0.75/IND	2	0.5	2/0.063	0.62/IND
VPCI 487/P/14 <sup>a</sup>	1	≥64	0.5/8	0.56/IND	1	1	0.5/0.125	0.62/IND
VPCI 518/P/14 <sup>a</sup>	0.5	≥64	0.25/8	0.56/IND	0.5	1	0.25/0.25	0.75/IND

<sup>a</sup>Fluconazole-resistant isolates (*n* = 10).<sup>b</sup>CAS, caspofungin; FLU, fluconazole; VRC, voriconazole; FICI, fractional inhibitory concentration index; IND, indifference; SYN, synergy; INT, interpretation.

and ≥8 μg/ml for fluconazole- and echinocandin-resistant species, respectively) (Tables 1 and 2) (14). All isolates were previously identified by conventional and molecular methods, i.e., CHROMagar *Candida* medium (Difco, Becton Dickinson & Company, Baltimore, MD, USA), microscopic morphology on cornmeal agar (Difco Laboratories, Detroit, MI, USA) with 1% Tween 80, and sequencing of internal transcribed spacer ribosomal DNA (rDNA) and D1/D2 regions. In addition, the isolates were identified by matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI Biotyper OC version 3.1; Bruker Daltonics, Bremen, Germany) (18). All strains were stored in 10% glycerol broth at –80°C at the Department of Medical Mycology, Vallabhbai Patel Chest Institute, University of Delhi, and were subcultured on Sabouraud dextrose agar (SDA) supplemented with 0.02% chloramphenicol at 35°C for 3 days to ensure purity and viability. All isolates were subcultured again on SDA before preparation of the inoculum. The interactions of caspofungin and micafungin with fluconazole or voriconazole were investigated by using a microdilution checkerboard method based on the CLSI reference technique with 96-well microtiter plates (24). Fluconazole (Pfizer, Groton, CT, USA), voriconazole (Pfizer), caspofungin (Merck), and micafungin (Astellas, Toyama, Japan) were dissolved in 100% dimethyl sulfoxide (DMSO). Drug dilutions were prepared to obtain four times the final concentration. Concentrations ranged from 8 to 0.016 μg/ml for caspofungin, 8 to 0.016 and 1 to 0.002 μg/ml for micafungin, 64 to 1 μg/ml for fluconazole, and 16 to 0.25 and 1 to 0.016

**TABLE 2** *In vitro* interactions of micafungin with fluconazole and voriconazole against *Candida auris*

Strain no.	MFG + FLU <sup>c</sup>				MFG + VRC <sup>c</sup>			
	MIC (μg/ml)				MIC (μg/ml)			
	MFG	FLU	MFG/FLU	FICI/INT	MFG	VRC	MFG/VRC	FICI/INT
VPCI 482/P/13 <sup>a</sup>	0.25	≥64	0.25/64	1.5/IND	0.25	2	0.016/0.5	0.31/SYN
VPCI 1132/P/13 <sup>a</sup>	0.5	32	0.25/4	0.62/IND	0.5	0.5	0.016/0.125	0.28/SYN
VPCI 1133/P/13 <sup>a,b</sup>	8	≥64	4/32	0.75/IND	8	1	2/0.25	0.5/SYN
VPCI 265/P/14 <sup>a</sup>	0.5	32	0.5/8	1.25/IND	0.5	8	0.063/1	0.25/SYN
VPCI 1510/P/14 <sup>a</sup>	0.125	32	0.063/8	0.75/IND	0.125	4	0.016/0.25	0.19/SYN
VPCI 1514/P/14 <sup>a,b</sup>	8	≥64	8/16	1.12/IND	8	0.5	1/0.125	0.37/SYN
VPCI 266/P/14 <sup>a</sup>	0.25	≥64	0.25/32	1.25/IND	0.25	0.5	0.008/0.125	0.28/SYN
VPCI 267/P/14 <sup>a,b</sup>	8	32	8/8	1.25/IND	8	0.5	1/0.125	0.37/SYN
VPCI 487/P/14 <sup>a</sup>	4	≥64	4/32	1.25/IND	4	1	0.5/0.125	0.25/SYN
VPCI 518/P/14 <sup>a</sup>	0.5	≥64	0.25/64	1/IND	0.5	1	0.016/0.125	0.15/SYN

<sup>a</sup>Fluconazole-resistant isolates (*n* = 10).<sup>b</sup>Micafungin-resistant isolates (*n* = 3).<sup>c</sup>MFG, micafungin; FLU, fluconazole; VRC, voriconazole; FICI, fractional inhibitory concentration index; IND, indifference; SYN, synergy; INT, interpretation.

$\mu\text{g/ml}$  for voriconazole. The concentration ranges of micafungin and voriconazole depended on the MIC results of each isolate. For two-dimensional microplate preparation, i.e., caspofungin plus fluconazole, caspofungin plus voriconazole, micafungin plus fluconazole, and micafungin plus voriconazole, 50  $\mu\text{l}$  of each concentration of echinocandins (caspofungin and micafungin) was added to columns 1 through 11, and then 50  $\mu\text{l}$  of azoles (fluconazole and voriconazole) was added to rows A through H, respectively. The wells of column 11 and the wells of row H contained 50  $\mu\text{l}$  of RPMI medium containing 1% of the solvent. Row H and column 11 contained the echinocandins and azoles alone, respectively. Column 12 was the drug-free well that served as the growth control. The maximal final concentration of DMSO in the test wells was  $<1\%$ . Trays were stored at  $-80^\circ\text{C}$  until the day of testing. After the microtiter trays were defrosted, 100  $\mu\text{l}$  of the inoculum was added to each well. Briefly, homogeneous suspensions were measured spectrophotometrically at 530 nm wavelength to a percentage transmission in the range of 75% to 77%. The final concentration of the stock inoculum suspensions of the isolates tested ranged from 1 to  $3 \times 10^3$  CFU/ml, as determined by quantitative colony counts on Sabouraud glucose agar (Difco). Plates were incubated at  $35^\circ\text{C}$  and examined visually after 24 h to determine MIC values for the drugs alone and in combination. The MIC endpoints were determined with the aid of a reading mirror and were defined as the lowest concentration of drug that significantly reduced growth ( $\geq 50\%$ ) compared with the growth of a drug-free control. For calculations, high off-scale MICs were raised to the next  $\log_2$  dilution step, while the low off-scale MICs were left unchanged (25). To assess the interactions of combinations of drugs, we calculated the fractional inhibitory concentration index (FICI). The FICI was defined as  $\text{FICI} = \text{FIC}_A + \text{FIC}_B = (C_A/\text{MIC}_A) + (C_B/\text{MIC}_B)$ , where  $\text{MIC}_A$  and  $\text{MIC}_B$  are the MICs of drugs A and B alone, and  $C_A$  and  $C_B$  are the concentrations of the drugs in combination, in all wells corresponding to an MIC. The interaction was considered synergistic when the FICI was  $\leq 0.5$ , indifferent at  $>0.5$  to  $\leq 4.0$ , and antagonistic at  $>4$  (24).

The results for the tested drugs alone and in combination against 10 *C. auris* strains are summarized in Tables 1 and 2. The MIC ranges of drugs alone against the strains were 32 to  $\geq 64$   $\mu\text{g/ml}$  for fluconazole, 0.5 to 8  $\mu\text{g/ml}$  for voriconazole, 0.5 to 4  $\mu\text{g/ml}$  for caspofungin, and 0.125 to 8  $\mu\text{g/ml}$  for micafungin. Based on findings with the checkerboard microdilution assay, when caspofungin was combined with fluconazole, the MIC ranges for caspofungin and fluconazole decreased to 0.25 to 2  $\mu\text{g/ml}$  and 8 to 64  $\mu\text{g/ml}$ , respectively; the combination exhibited indifferent activity against all 10 strains (FICI, 0.56 to 2). When caspofungin was combined with voriconazole, the MIC ranges for caspofungin and voriconazole decreased to 0.25 to 2  $\mu\text{g/ml}$  and 0.063 to 4  $\mu\text{g/ml}$ , respectively, and demonstrated indifferent activity against all strains (FICI, 0.62 to 2) (Table 1). When micafungin was combined with fluconazole, the MIC ranges of micafungin and fluconazole were reduced to 0.063 to 8  $\mu\text{g/ml}$  and 4 to 64  $\mu\text{g/ml}$ , respectively; indifference was also observed (FICI, 0.62 to 1.5) (Table 2). Synergistic effects of micafungin with voriconazole were shown against the 10 multidrug-resistant *C. auris* isolates (FICI, 0.15 to 0.5); the MIC ranges of micafungin and voriconazole were reduced to 0.008 to 2  $\mu\text{g/ml}$  and 0.125 to 1  $\mu\text{g/ml}$ , respectively (Table 2). Overall, no antagonistic effects were observed for any combination.

In this study, we used the checkerboard microdilution method to analyze drug-drug interactions of echinocandins with azoles against multidrug-resistant *C. auris*. The emergence of new species and antifungal resistance has raised the issue of using alternative therapeutic strategies. Evidence to support treatment choices for multidrug-resistant *C. auris* disease is rare. Except for one study (20), *in vitro* antifungal profiles are relatively scarce and based on low numbers of test isolates (14, 19, 21). The *in vivo* efficacy of antifungal therapy against *C. auris* is undetermined, and *in vitro* data from different sources are inadequate. Use of echinocandins is the recommended treatment for patients with potent activity, an excellent safety profile, and favorable pharmacokinetics (26–28), but unsuccessful treatment of *C. auris* infections with fluconazole, voriconazole, amphotericin B, caspofungin, and anidulafungin has been reported (6).

On the other hand, micafungin is used for prophylaxis and treatment with a broad spectrum of activity in both neutropenic and nonneutropenic patients (15, 29). Concordant with other reports (30–32), micafungin activity was shown to be as effective as caspofungin *in vitro* against *Candida glabrata* isolates with and without *fks* mutations. Micafungin was also effective *in vivo* for decreasing the fungal burden in mice infected with *C. glabrata* with *fks* mutations. It seems that lower concentrations of drugs cause fewer side effects and improve the treatment outcomes. We have shown that interaction between micafungin and voriconazole exhibited synergistic activity against multidrug-resistant *C. auris* strains, suggesting that the combination may be considered for patients with candidiasis. However, *in vivo* studies with suitable animal models of *C. auris* infection are needed to confirm the *in vitro* results presented here. Clearly, more research is indicated to explore clinical management. In conclusion, the combination of micafungin and voriconazole exhibited synergistic activity against multidrug-resistant *C. auris*, suggesting that this is an alternative approach to overcome antifungal drug resistance. However, use of this combination therapy *in vivo* and determination of the underlying mechanism of this synergistic action need further study.

### ACKNOWLEDGMENTS

This study was financially supported by a grant from the Department of Medical Mycology, Vallabhbhai Patel Chest Institute, University of Delhi, Delhi, India. H.F. was supported by an ESCMID observership grant (ID 1002), which is gratefully acknowledged. H.B. was funded by a grant (no. 2477) from Mazandaran University of Medical Sciences, Sari, Iran.

J.F.M. received grants from Astellas, Merck, and Basilea. He has been a consultant to Basilea and Merck and received speaker fees from Merck, Pfizer, Gilead, and United Medical. During the past 5 years, E.D. has received research grants from MSD and Gilead; travel grants from Gilead, MSD, Pfizer, and Astellas; and speaker fees from Gilead, MSD, and Astellas. All other authors have no potential conflicts of interest.

We alone are responsible for the content and writing of the paper.

### REFERENCES

- Pfaller MA, Andes DR, Diekema DJ, Horn DL, Reboli AC, Rotstein C, Franks B, Azie NE. 2014. Epidemiology and outcomes of invasive candidiasis due to non-*albicans* species of *Candida* in 2,496 patients: data from the prospective Antifungal Therapy (PATH) registry 2004–2008. *PLoS One* 9:e101510. <https://doi.org/10.1371/journal.pone.0101510>.
- Chowdhary A, Sharma C, Meis JF. 2017. *Candida auris*: a rapidly emerging cause of hospital-acquired multidrug-resistant fungal infections globally. *PLoS Pathog* 13:e1006290. <https://doi.org/10.1371/journal.ppat.1006290>.
- Rudramurthy SM, Chakrabarti A, Paul RA, Sood P, Kaur H, Kapoor MR, Kindo AJ, Marak RS, Arora A, Sardana R, Das S, Chhina D, Patel A, Xess I, Tarai B, Singh P, Ghosh A. 2017. *Candida auris* candidaemia in Indian ICUs: analysis of risk factors. *J Antimicrob Chemother* 72:1794–1801. <https://doi.org/10.1093/jac/dkx034>.
- Chowdhary A, Sharma C, Duggal S, Agarwal K, Prakash A, Singh PK, Jain S, Kathuria S, Randhawa HS, Hagen F, Meis JF. 2013. New clonal strain of *Candida auris*, Delhi, India. *Emerg Infect Dis* 19:1670–1673. <https://doi.org/10.3201/eid1910.130393>.
- Chowdhary A, Kumar VA, Sharma C, Prakash A, Agarwal K, Babu R, Dinesh K, Karim S, Singh S, Hagen F. 2014. Multidrug-resistant endemic clonal strain of *Candida auris* in India. *Eur J Clin Microbiol Infect Dis* 33:919–926. <https://doi.org/10.1007/s10096-013-2027-1>.
- Calvo B, Melo AS, Perozo-Mena A, Hernandez M, Francisco EC, Hagen F, Meis JF, Colombo AL. 2016. First report of *Candida auris* in America: clinical and microbiological aspects of 18 episodes of candidemia. *J Infect* 73:369–374. <https://doi.org/10.1016/j.jinf.2016.07.008>.
- Mohsin J, Hagen F, Al-Balushi ZAM, de Hoog GS, Chowdhary A, Meis JF, Al-Hatmi AMS. 2017. The first cases of *Candida auris* candidaemia in Oman. *Mycoses* 60:569–575. <https://doi.org/10.1111/myc.12647>.
- Magobo RE, Corcoran C, Seetharam S, Govender NP. 2014. *Candida auris*–associated candidemia, South Africa. *Emerg Infect Dis* 20:1250–1251.
- Schelenz S, Hagen F, Rhodes JL, Abdolrasouli A, Chowdhary A, Hall A, Ryan L, Shackleton J, Trimlett R, Meis JF, Armstrong-James D, Fisher MC. 2016. First hospital outbreak of the globally emerging *Candida auris* in a European hospital. *Antimicrob Resist Infect Control* 5:35. <https://doi.org/10.1186/s13756-016-0132-5>.
- Borman AM, Szekeley A, Johnson EM. 2017. Isolates of the emerging pathogen *Candida auris* present in the UK have several geographic origins. *Med Mycol* 55:563–567. <https://doi.org/10.1093/mmy/myw147>.
- Ben-Ami R, Berman J, Novikov A, Bash E, Shachor-Meyouhas Y, Zakin S, Maor Y, Tarabia J, Schechner V, Adler A. 2017. Multidrug-resistant *Candida haemulonii* and *C. auris*, Tel Aviv, Israel. *Emerg Infect Dis* 23:195–203.
- Vallabhaneni S, Kallen A, Tsay S, Chow N, Welsh R, Kerins J, Kembler SK, Pacilli M, Black SR, Landon E, Ridgway J, Palmore TN, Zelzany A, Adams EH, Quinn M, Chaturvedi S, Greenko J, Fernandez R, Southwick K, Furuya EY, Calfee DP, Hamula C, Patel G, Barrett P, Lafaro P, Berkow EL, Moulton-Meissner H, Noble-Wang J, Fagan RP, Jackson BR, Lockhart SR, Litvintseva AP, Chiller TM. 2017. Investigation of the first seven reported cases of *Candida auris*, a globally emerging invasive, multidrug-resistant fungus–United States, May 2013–August 2016. *Am J Transplant* 17:296–299.
- Morales-López SE, Parra-Giraldo CM, Ceballos-Garzón A, Martínez HP, Rodríguez GJ, Álvarez-Moreno CA, Rodríguez JY. 2017. Invasive infections with multidrug-resistant yeast *Candida auris*, Colombia. *Emerg Infect Dis* 23:162–164. <https://doi.org/10.3201/eid2301.161497>.
- Lockhart SR, Etienne KA, Vallabhaneni S, Farooqi J, Chowdhary A, Govender NP, Colombo AL, Calvo B, Cuomo CA, Desjardins CA, Berkow EL, Castanheira M, Magobo RE, Jabeen K, Asghar RJ, Meis JF, Jackson B,

- Chiller T, Litvintseva AP. 2017. Simultaneous emergence of multidrug-resistant *Candida auris* on 3 continents confirmed by whole-genome sequencing and epidemiological analyses. *Clin Infect Dis* 64:134–140. <https://doi.org/10.1093/cid/ciw691>.
15. Cornely OA, Bassetti M, Calandra T, Garbino J, Kullberg B, Lortholary O, Meersseman W, Akova M, Arendrup M, Arikan-Akdagli S, Bille J, Castagnola E, Cuenca-Estrella M, Donnelly JP, Groll AH, Herbrecht R, Hope WW, Jensen HE, Lass-Flörl C, Petrikos G, Richardson MD, Roilides E, Verweij PE, Viscoli C, Ullmann AJ. 2012. ESCMID guideline for the diagnosis and management of *Candida* diseases 2012: non-neutropenic adult patients. *Clin Microbiol Infect* 18:19–37. <https://doi.org/10.1111/1469-0691.12039>.
  16. Arendrup MC, Boekhout T, Akova M, Meis JF, Cornely OA, Lortholary O. 2014. ESCMID and ECMM joint clinical guidelines for the diagnosis and management of rare invasive yeast infections. *Clin Microbiol Infect* 20(Suppl 3):S76–S98. <https://doi.org/10.1111/1469-0691.12360>.
  17. Gonçalves SS, Souza AC, Chowdhary A, Meis JF, Colombo AL. 2016. Epidemiology and molecular mechanisms of antifungal resistance in *Candida* and *Aspergillus*. *Mycoses* 59:198–219. <https://doi.org/10.1111/myc.12469>.
  18. Kathuria S, Singh PK, Sharma C, Prakash A, Masih A, Kumar A, Meis JF, Chowdhary A. 2015. Multidrug-resistant *Candida auris* misidentified as *Candida haemulonii*: characterization by matrix-assisted laser desorption ionization–time of flight mass spectrometry and DNA sequencing and its antifungal susceptibility profile variability by Vitek 2, CLSI broth microdilution, and Etest method. *J Clin Microbiol* 53:1823–1830. <https://doi.org/10.1128/JCM.00367-15>.
  19. Larkin E, Hager C, Chandra J, Mukherjee PK, Retuerto M, Salem I, Long L, Isham N, Kovanda L, Borroto-Esoda K, Wring S, Angulo D, Ghannoum M. 2017. The emerging pathogen *Candida auris*: growth phenotype, virulence factors, activity of antifungals, and effect of SCY-078, a novel glucan synthesis inhibitor, on growth morphology and biofilm formation. *Antimicrob Agents Chemother* 61:pii=e02396-16. <https://doi.org/10.1128/AAC.02396-16>.
  20. Arendrup MC, Prakash A, Meletiadis J, Sharma C, Chowdhary A. 2017. *Candida auris*: comparison of the EUCAST and CLSI reference microdilution MICs for eight antifungal compounds and associated tentative ECOFFs. *Antimicrob Agents Chemother* 61:pii=e00485-17. <https://doi.org/10.1128/AAC.00485-17>.
  21. Berkow EL, Angulo D, Lockhart SR. 2017. *In vitro* activity of a novel glucan synthase inhibitor, SCY-078, against clinical isolates of *Candida auris*. *Antimicrob Agents Chemother* 61:pii=e00435-17. <https://doi.org/10.1128/AAC.00435-17>.
  22. Mizusawa M, Miller H, Green R, Lee R, Durante M, Perkins R, Hewitt C, Simner PJ, Carroll KC, Hayden RT, Zhang SX. 2017. Can multidrug-resistant *Candida auris* be reliably identified in clinical microbiology laboratories? *J Clin Microbiol* 55:638–640. <https://doi.org/10.1128/JCM.02202-16>.
  23. Fakhim H, Emami S, Vaezi A, Hashemi SM, Faeli L, Diba K, Dannaoui E, Badali H. 2016. *In vitro* activities of novel azole compounds (ATTAF-1 and ATTAF-2) against fluconazole-susceptible and -resistant isolates of *Candida* species. *Antimicrob Agents Chemother* 61:pii=e01106-16. <https://doi.org/10.1128/AAC.01106-16>.
  24. Odds FC. 2003. Synergy, antagonism, and what the checkerboard puts between them. *J Antimicrob Chemother* 52:1. <https://doi.org/10.1093/jac/dkg301>.
  25. Pfaller MA, Watanabe N, Castanheira M, Messer SA, Jones RN. 2011. Pre-clinical development of antifungal susceptibility test methods for the testing of the novel antifungal agent E1210 versus *Candida*: comparison of CLSI and European Committee on Antimicrobial Susceptibility Testing methods. *J Antimicrob Chemother* 66:2581–2584. <https://doi.org/10.1093/jac/dkr342>.
  26. Pfaller M, Boyken L, Hollis R, Kroeger J, Messer S, Tendolkar S, Diekema D. 2008. *In vitro* susceptibility of invasive isolates of *Candida* spp. to anidulafungin, caspofungin, and micafungin: six years of global surveillance. *J Clin Microbiol* 46:150–156. <https://doi.org/10.1128/JCM.01901-07>.
  27. Dannaoui E, Lortholary O, Raoux D, Bougnoux M, Galeazzi G, Lawrence C, Moissenet D, Poilane I, Hoinard D, Dromer F. 2008. Comparative *in vitro* activities of caspofungin and micafungin, determined using the method of the European Committee on Antimicrobial Susceptibility Testing, against yeast isolates obtained in France in 2005–2006. *Antimicrob Agents Chemother* 52:778–781. <https://doi.org/10.1128/AAC.01140-07>.
  28. Prigent G, Ait-Ammar N, Levesque E, Fekkar A, Costa J-M, El Abbassi S, Foulet F, Duvoux C, Merle J-C, Dannaoui E. 2017. Echinocandin resistance in *Candida* spp. isolated from liver transplant recipients. *Antimicrob Agents Chemother* 61:e01229-16. <https://doi.org/10.1128/AAC.01229-16>.
  29. Pfaller MA, Messer SA, Woosley LN, Jones RN, Castanheira M. 2013. Echinocandin and triazole antifungal susceptibility profiles of opportunistic yeast and mould clinical isolates (2010–2011): application of new CLSI clinical breakpoints and epidemiological cutoff values to characterize geographic and temporal trends of antifungal resistance. *J Clin Microbiol* 51:2571–2581. <https://doi.org/10.1128/JCM.00308-13>.
  30. Lepak A, Castanheira M, Diekema D, Pfaller M, Andes D. 2012. Optimizing echinocandin dosing and susceptibility breakpoint determination via *in vivo* pharmacodynamic evaluation against *Candida glabrata* with and without *FKS* mutations. *Antimicrob Agents Chemother* 56:5875–5882. <https://doi.org/10.1128/AAC.01102-12>.
  31. Arendrup MC, Perlin DS, Jensen RH, Howard SJ, Goodwin J, Hope W. 2012. Differential *in vivo* activities of anidulafungin, caspofungin, and micafungin against *Candida glabrata* isolates with and without *FKS* resistance mutations. *Antimicrob Agents Chemother* 56:2435–2442. <https://doi.org/10.1128/AAC.06369-11>.
  32. Spreghini E, Orlando F, Sanguinetti M, Posteraro B, Giannini D, Manso E, Barchiesi F. 2012. Comparative effects of micafungin, caspofungin, and anidulafungin against a difficult-to-treat fungal opportunistic pathogen, *Candida glabrata*. *Antimicrob Agents Chemother* 56:1215–1222. <https://doi.org/10.1128/AAC.05872-11>.